


### Descomposición de un paralelepípedo en pirámides de base cuadrilátera

Define el paralelepípedo...

$$\begin{aligned}
 A & \left( \begin{array}{c} \uparrow \\ \downarrow \end{array} -0.50, \begin{array}{c} \uparrow \\ \downarrow \end{array} -0.50, \begin{array}{c} \uparrow \\ \downarrow \end{array} 0.50 \right) & D & \left( \begin{array}{c} \uparrow \\ \downarrow \end{array} -0.50, \begin{array}{c} \uparrow \\ \downarrow \end{array} 0.50, \begin{array}{c} \uparrow \\ \downarrow \end{array} 0.75 \right) \\
 B & \left( \begin{array}{c} \uparrow \\ \downarrow \end{array} 0.50, \begin{array}{c} \uparrow \\ \downarrow \end{array} -0.50, \begin{array}{c} \uparrow \\ \downarrow \end{array} 0.50 \right) & E & \left( \begin{array}{c} \uparrow \\ \downarrow \end{array} -0.50, \begin{array}{c} \uparrow \\ \downarrow \end{array} -0.50, \begin{array}{c} \uparrow \\ \downarrow \end{array} -0.50 \right)
 \end{aligned}$$



Romboiedro

Indicaciones

Autor: José R. Galo Sánchez

Proyecto Descartes. Año 2018